



CLAIMS

1. A method for preparing a polyurethane foam which comprises reacting an organic polyisocyanate and a polyol in the presence of water as a blowing agent, a cell stabilizer, and a tertiary amino alkyl amide catalyst composition represented by the formula I:



wherein A represents CH or N;

10 R¹ represents hydrogen or




n is an integer from 1 to 3;

15 R² and R³ each represent hydrogen or a C₁-C₆ linear or branched alkyl group;

R⁴ and R⁵ each represent a C₁-C₆ linear or branched alkyl group when A represents N, or together R⁴ and R⁵ represent a C₂-C₅ alkylene group when A represents N; or together R⁴ and R⁵ may be a C₂-C₅ alkylene group containing NR⁷ when A represents CH or N, where R⁷ is selected from the group consisting of hydrogen, a C₁-C₄ linear or branched

20 alkyl group, and



and; R⁶ represents a C₅-C₃₅ linear or branched alkyl, alkenyl, or aryl group, and where the tertiary amino alkyl amide catalyst is acid-blocked.

25

2. The method of claim 1, wherein R^1 , R^2 , and R^3 each represent hydrogen.

3. The method of claim 1, wherein R^4 and R^5 each represent a methyl group
5 when A represents N.

4. The method of claim 1, wherein R^4 and R^5 together represent
-CH₂CH₂N(CH₃)CH₂- when A represents CH.

10 5. The method of claim 1, wherein n represents 2 or 3.

6. The method of claim 1, wherein the tertiary amino alkyl amide catalyst
composition is an N-(3-dimethylaminopropyl)-amide derived from an acid selected from
the group consisting of 2-ethylhexanoic, coconut oil fatty, tall oil fatty, caproic, heptylic,
15 caprylic, pelargonic, capric, hendecanoic, lauric, tridecanoic, myristic, pentadecanoic,
palmitic, margaric, stearic, oleic, linoleic, linolenic, ricinoleic, nonadecanoic, arachidic,
heneicosanoic, behenic, tricosanoic, lignoceric, pentacosanoic, cerotic, heptacosanoic,
montanic, nonacosanoic, melissic, hentriacontanoic, dotriacontanoic, tritriacontanoic,
tetracontanoic, hexatriacontanoic, 9-phenylstearic, and 10-phenylstearic acid.

20

7. The method of claim 6, wherein the acid is selected from the group consisting
of 2-ethylhexanoic, coconut oil fatty, and tall oil fatty acids.

8. The method of claim 1, wherein the tertiary amino alkyl amide catalyst
25 composition is an N-(2-dimethylaminoethyl)-amide derived from an acid selected from
the group consisting of 2-ethylhexanoic, coconut oil fatty, tall oil fatty, caproic, heptylic,
caprylic, pelargonic, capric, hendecanoic, lauric, tridecanoic, myristic, pentadecanoic,
palmitic, margaric, stearic, oleic, linoleic, linolenic, ricinoleic, nonadecanoic, arachidic,
heneicosanoic, behenic, tricosanoic, lignoceric, pentacosanoic, cerotic, heptacosanoic,
30 montanic, nonacosanoic, melissic, hentriacontanoic, dotriacontanoic, tritriacontanoic,
tetracontanoic, hexatriacontanoic, 9-phenylstearic, and 10-phenylstearic acid.

9. The method of claim 8, wherein the acid is selected from the group consisting
of 2-ethylhexanoic, coconut oil fatty, and tall oil fatty acids.

10. The method of claim 1, wherein the tertiary amino alkyl amide catalyst composition is an N-methyl-3-aminoethyl pyrrolidine amide derived from an acid selected from the group consisting of 2-ethylhexanoic, coconut oil fatty, tall oil fatty, caproic, 5 heptylic, caprylic, pelargonic, capric, hendecanoic, lauric, tridecanoic, myristic, pentadecanoic, palmitic, margaric, stearic, oleic, linoleic, linolenic, ricinoleic, nonadecanoic, arachidic, heneicosanoic, behenic, tricosanoic, lignoceric, pentacosanoic, cerotic, heptacosanoic, montanic, nonacosanoic, melissic, hentriacontanoic, dotriacontanoic, tritriacontanoic, tetracontanoic, hexatriacontanoic, 9-phenylstearic, and 10 10-phenylstearic acid.

11. The method of claim 10, wherein the acid is selected from the group consisting of 2-ethylhexanoic, coconut oil fatty, and tall oil fatty acids.

15 12. The method of claim 1, wherein the tertiary amino alkyl amide catalyst composition is a 4, 10-diaza-4, 10, 10-trimethyl-7-oxa-undecaamine amide derived from an acid selected from the group consisting of 2-ethylhexanoic, coconut oil fatty, tall oil fatty, caproic, heptylic, caprylic, pelargonic, capric, hendecanoic, lauric, tridecanoic, myristic, pentadecanoic, palmitic, margaric, stearic, oleic, linoleic, linolenic, ricinoleic, 20 nonadecanoic, arachidic, heneicosanoic, behenic, tricosanoic, lignoceric, pentacosanoic, cerotic, heptacosanoic, montanic, nonacosanoic, melissic, hentriacontanoic, dotriacontanoic, tritriacontanoic, tetracontanoic, hexatriacontanoic, 9-phenylstearic, and 10-phenylstearic acid.

25 13. The method of claim 12, wherein the acid is selected from the group consisting of 2-ethylhexanoic, coconut oil fatty, and tall oil fatty acids.

14. The method of claim 1, wherein the tertiary amino alkyl amide catalyst composition is N-(3-dimethylaminopropyl)-2-ethyl-hexamide.

30 15. The method of claim 1, wherein the tertiary amino alkyl amide catalyst composition is N-(3-dimethylaminopropyl)-cocoamide.

16. The method of claim 1, wherein the tertiary amino alkyl amide catalyst composition is N-(3-dimethylaminopropyl)-tall oil amide.


17. The method of claim 1, which comprises reacting the following components
5 in parts by weight (pbw):

Polyol	20-100
Polymer Polyol	80-0
Silicone Surfactant	1-2.5
Blowing agent	2-4.5
Crosslinker	0.5-2
Catalyst	0.25-2
Isocyanate Index	70-115


18. The method of Claim 1 in which the catalyst composition is acid-blocked with a carboxylic acid.

19. The method of Claim 18 in which the carboxylic acid is formic acid, acetic acid, 2-ethyl-hexanoic acid, gluconic acid, or N-(2-hydroxyethyl)-iminodiacetic acid.

20. In a method for preparing a polyurethane foam which comprises reacting an
15 organic polyisocyanate and a polyol in the presence of water as a blowing agent, a cell
stabilizer, and a catalyst composition, the improvement of controlling and improving the
porosity and openness of the foam which comprises using a tertiary amino alkyl amide
catalyst composition represented by the formula I:



wherein A represents CH or N;
R¹ represents hydrogen or




n is an integer from 1 to 3;

R² and R³ each represent hydrogen or a C₁-C₆ linear or branched alkyl group;

- 5 R⁴ and R⁵ each represent a C₁-C₆ linear or branched alkyl group when A represents N, or together R⁴ and R⁵ represent a C₂-C₅ alkylene group when A represents N; or together R⁴ and R⁵ may be a C₂-C₅ alkylene group containing NR⁷ when A represents CH or N, where R⁷ is selected from the group consisting of hydrogen, a C₁-C₄ linear or branched alkyl group, and

10



and; R⁶ represents a C₅-C₃₅ linear or branched alkyl, alkenyl, or aryl group, and where the tertiary amino alkyl amide catalyst is acid-blocked with a carboxylic acid.

15

ML03167(FINAL).APP.DOC

20